

## **SELF MANAGEMENT STRATEGIES OF OLDER PEOPLE WITH ASTHMA**

D Goeman<sup>1</sup>, R O'Hehir<sup>1</sup>, S Scharf<sup>3</sup>, C Jenkins<sup>2</sup> J Douglass<sup>1</sup>.

<sup>1</sup> CRC for Asthma and Airways, The Alfred Hospital, Melbourne, Victoria; <sup>2</sup> CRC for Asthma and Airways, Woolcock Institute, Sydney, NSW, <sup>3</sup> Aged Care Services, Caulfield General Medical Centre, Caulfield, Victoria

**Purpose:** To obtain the perspectives of older people with asthma in order to understand the reasons for the higher risk of death from asthma in this group.

**Methods:** Combined qualitative and quantitative study of people over 50 years of age. Fifty five participants (16 male, 39 female) were recruited from inner city & suburban areas, predominantly surrounding a tertiary hospital, and the regional area of South Gippsland, an area with a high population of retirees. Participants were interviewed in-depth on their experiences, attitudes and beliefs about asthma. Lung function was measured pre and post beta 2 agonist use and questionnaire data on background information, asthma symptoms, asthma knowledge, asthma control, medication use and respiratory health was also collected.

**Results:** Findings from this study revealed barriers to the self-management of older people with asthma and in the provision of asthma care by health professionals. Education on asthma treatments and self-management strategies needs to be tailored in accordance with the age and time of asthma onset and the duration of diagnosis.

**Conclusions:** In order to improve asthma outcomes in older people we believe a specifically targeted intervention is needed to improve the content and quality of consultations for older people with asthma.

Supported by: Asthma Victoria and Co-operative Research Centre for Asthma and Airways